

CONVENIA® (cefovecin sodium) in your practice

- Label dose for dogs and cats = 3.6 mg/lb (8 mg/kg or 0.045 mL/lb) subcutaneously
- Reconstituted concentration = 80 mg/mL
- Reconstitute with 10 mL of sterile water for injection, USP
- Use within 56 days after reconstitution
- Always refrigerate in original carton to protect from light
- Minimum pet age for use is 4 months

As with other cephalosporins, CONVENIA will naturally turn amber over time after reconstitution. The color change has no impact on safety or efficacy, provided CONVENIA is kept under proper storage conditions.



Examples of color change after the vial has been broached and reconstituted.

CONVENIA Dosing Chart at 3.6 mg/lb (8 mg/kg or 0.045 mL/lb)

Weight of Animal	Volume of CONVENIA
5 lb	0.23 mL
10 lb	0.45 mL
15 lb	0.67 mL
20 lb	0.90 mL
40 lb	1.80 mL
60 lb	2.70 mL
80 lb	3.60 mL
100 lb	4.50 mL

Important Safety Information:

People with known hypersensitivity to penicillin or cephalosporins should avoid exposure to CONVENIA. Do not use in dogs or cats with a history of allergic reactions to penicillins or cephalosporins. Side effects for both dogs and cats include vomiting, diarrhea, decreased appetite/anorexia and lethargy. For more safety information, please see accompanying Full Prescribing Information.

convenia
cefovecin sodium

zoetis