

## Chlortetracycline Beef Cattle Feed – AUREO<sup>a</sup> – BE-BP – TD

### Type B Medicated Feed - TD

(chlortetracycline Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For calves, beef and non-lactating dairy cattle: Treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to chlortetracycline.

#### Active Drug Ingredient

Chlortetracycline<sup>a</sup> ..... 20,001 – 80,000 g/ton\*

#### Guaranteed Analysis

Crude Protein (Min) .....	_____	%
NPN <sup>1</sup> (Max) .....	_____	%
Crude Fat (Min) .....	_____	%
Crude Fiber (Max) .....	_____	%
Calcium <sup>1</sup> (Min) .....	_____	%
Calcium <sup>1</sup> (Max) .....	_____	%
Phosphorus <sup>1</sup> (Min) .....	_____	%
Salt <sup>1</sup> (Min) .....	_____	%
Salt <sup>1</sup> (Max) .....	_____	%
Sodium <sup>2</sup> (Min) .....	_____	%
Sodium <sup>2</sup> (Max) .....	_____	%
Potassium <sup>1</sup> (Min) .....	_____	%
Vitamin A <sup>1</sup> .....	_____	IU/lb

<sup>1</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>2</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

#### Ingredients

Ingredients as defined by AAFCO.

#### Mixing Directions

Mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C top-dress medicated feed for cattle containing 4,000 – 20,000 g chlortetracycline per ton.

*The following table provides examples of mixing rates:*

Type B CTC concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C CTC concentration (g/ton)
40,000 (20 g/lb)	200	1,800	4,000
80,000 (40 g/lb)	100	1,900	4,000
80,000 (40 g/lb)	500	1,500	20,000

The resulting Type C medicated feed should be fed for not more than 5 days to provide chlortetracycline at the rate of 10 mg/pound of body weight per day. The medicated feed should be administered as a top dress or mixed into the daily ration. If top-dressed, spread uniformly on top of ration and allow sufficient space so that all cattle can eat at the same time.

**Warning**

A withdrawal period has not been established for this product in pre-ruminating calves.  
Do not use in calves to be processed for veal.

Lot Number (if applicable): \_\_\_\_\_

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

\* The final printed medicated feed label must state a single drug concentration.

<sup>a</sup>Aureomycin® is the proprietary name of chlortetracycline Type A medicated article (NADA 048-761).

Zoetis Inc. [15 March 2016]