

Chlortetracycline-AUREO (10 mg/lb BW) + Laidlomycin (5-10 g/ton) CATTLE FEED
Type B Medicated Feed
(chlortetracycline and laidlomycin propionate potassium Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline, and for improved feed efficiency in cattle fed in confinement for slaughter.

ACTIVE DRUG INGREDIENTS*

Chlortetracycline^a 4000 to 80,000 g/ton
Laidlomycin propionate potassium^b 33.3 to 1200 g/ton

GUARANTEED ANALYSIS

Crude protein, not less than %
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %
Vitamin A¹ (not less than) IU/lb.

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

MIXING DIRECTIONS

In order to treat cattle for bacterial enteritis and/or pneumonia while improving feed efficiency, mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of complete cattle feed containing 500 to 4000 grams chlortetracycline and 5 to 10 grams laidlomycin propionate potassium. The following table provides examples of mixing rates.

Body Wt., lb.	Feed Consumption, lb./head/day	Type B Concentration, g/ton		Type B per Ton of Type C, lb.	Non-Medicated Feed per Ton of Type C, lb.	Type C Concentration, g/ton	
		Chlortetracycline	Laidlomycin			Chlortetracycline	Laidlomycin
500	8	10000.0	60.0	250	1750	1250.0	7.5
600	6	20000.0	100.0	200	1800	2000.0	10
600	12	6666.7	33.3	300	1700	1000.0	5
800	20	6000.0	37.5	266.7	1733.3	800.0	5
1000	6	12000.0	36.0	555.6	1444.4	3333.3	10
1000	30	66666.7	500.0	20.0	1980	666.7	5

The resulting Type C medicated feed should be fed continuously at a rate to provide 10 mg chlortetracycline per lb. body weight/day and 30 to 150 mg laidlomycin propionate potassium per head per day for not more than 5 days.

Caution

Do not allow horses or other equines access to feeds containing laidlomycin propionate potassium. The safety of laidlomycin propionate potassium in unapproved species has not been established. Not for use in animals intended for breeding.

Warning: No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.



Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

*Final printed label on formulated Type B Medicated feed must bear a single concentration for each drug.

^aAureomycin® is the proprietary name of chlortetracycline Type A medicated article (NADA 048-761).

^bCattlyst® is the proprietary name of laidlomycin propionate potassium Type A medicated article (NADA 141-025).

Zoetis Inc. [7 September 2016]