

Liquid Type C Medicated Cattle Feed M-T+
Type C Medicated Feed
(melengestrol acetate and tylosin Type C top-dress medicated feed)

For Use in Heifer Feeds Only

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS

Heifers being fed in confinement for slaughter: For increased rate of weight gain, improved feed efficiency, suppression of estrus (heat), and for reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*.

ACTIVE DRUG INGREDIENTS

Melengestrol acetate (MGA)^a.....0.0000276 to 0.00022% (0.25 to 2 g/ton)*
Tylosin (as tylosin phosphate) (TylanTM)^b.....90 to 360 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____ %
Non-Protein Nitrogen (NPN) ¹ , not more than	_____ %
Crude Fat, not less than.....	_____ %
Crude Fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than.....	_____ %
Salt ² , not less than.....	_____ %
Salt ² , not more than.....	_____ %
Sodium ³ , not less than.....	_____ %
Sodium ³ , not more than.....	_____ %
Potassium, not less than.....	_____ %
Vitamin A ^{2,4} , not less than.....	_____ I.U./lb
Dry Matter, not less than	60 %
Dry Matter, not more than.....	75%
pH.....	4.5 to 6.0

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

For stored liquid Type C medicated feeds containing melengestrol acetate and tylosin, recirculate or agitate liquid Type C medicated feeds daily even when no Type C feed is used and immediately prior to use for no less than 10 minutes moving not less than 1% of the tank contents per minute from the bottom to the top of the tank.

Each pound contains 0.125 to 1.0 mg melengestrol acetate and 45 to 180 mg of tylosin. Feed to heifers at a rate of 0.5 to 2.0 pounds per head per day to provide 0.25 to 0.5 mg melengestrol acetate and 60 to 90 mg tylosin per head per day. Prior to feeding, this Liquid Type C product must be top-dressed onto a complete feed or mixed into the amount of complete feed consumed by an animal per day.

Caution

Feed continuously as sole ration.

MGA is for use only in heifers being fed in confinement for slaughter. Not effective in steers and spayed heifers.

Warning: No withdrawal period is required.

When mixing and handling Tylan™, use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

Expiration Date: (31 days after manufacture)

*Final printed label on formulated Type C medicated feed must bear a single concentration for each drug.

^aMGA® is the proprietary name of melengestrol acetate Type A medicated article (NADA 039-402).

^bTylan™ is the proprietary name of tylosin phosphate Type A medicated article (NADA 012-491).

Zoetis Inc. [7 September 2016]