

**DCX + Chlortetracycline-AUREO Complete Cattle Feed Medicated
Type C Medicated Complete Feed**
(decoquinatate and chlortetracycline Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Calves, beef, and non-lactating dairy cattle: For the prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii*; for treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENTS*

Decoquinatate^a..... 12.9 to 90.8 g/ton*
Chlortetracycline^b (as hydrochloride)..... 500 to 4000 g/ton*

GUARANTEED ANALYSIS

Crude protein, not less than %
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

FEEDING DIRECTIONS

Feed at a rate of 1g chlortetracycline per 100 lb body weight/day and 22.7 mg decoquinatate per 100 lb of body weight/day for not more than 5 days. When it is fully consumed, resume feeding 22.7 mg decoquinatate per 100 lb of body weight/day for a total of 28 days to prevent coccidiosis. The following table provides examples of feeding rates and amount of drug per ton of feed.

Animal Wt. (lbs)	Feed Intake (% Body Wt)	Lbs Feed Per Head /Day	Decoquinatate (grams/ton)	Chlortetracycline, (grams/ton)
250	0.5	1.25	90.8	4000
	1.0	2.5	45.4	2000
	3.5	8.75	13.0	571
500	0.5	2.5	90.8	4000
	1.0	5.0	45.4	2000
	3.5	17.5	13.0	571
1000	0.5	5.0	90.8	4000
	1.0	10.0	45.4	2000
	3.5	35.0	13.0	571

Warning: No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to animals producing milk for food.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

*Final printed label on formulated Type C Medicated feed must bear a single concentration for each drug. The ratio between chlortetracycline and decoquinatate must be 44:1.

^aDecco^x® is the proprietary name of decoquinatate Type A medicated article (NADA 039-417).

^bAureomycin® is the proprietary name of chlortetracycline Type A medicated article (NADA 048-761).

Zoetis Inc. [9 November 2016]