

Chlortetracycline-AUREO (350 mg/head) + Laidlomycin (5 g/ton)
TYPE C Medicated Cattle Feed
(chlortetracycline and laidlomycin propionate potassium Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline, and for improved feed efficiency and increased rate of weight gain in cattle fed in confinement for slaughter.

ACTIVE DRUG INGREDIENTS*

Chlortetracycline^a 23.3 to 58.3 g/ton
Laidlomycin propionate potassium^b 5 g/ton

GUARANTEED ANALYSIS

Crude protein, not less than %
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %
Vitamin A¹ (not less than) IU/lb.

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

FEEDING DIRECTIONS

Provide this Type C medicated feed to confined beef cattle to control bacterial pneumonia while improving feed efficiency and increasing rate of weight gain. Feed continuously at a rate of 350 mg chlortetracycline per head per day and 30 to 75 mg laidlomycin propionate potassium per head per day. The following table provides examples of feeding rates and amount of drug per ton of feed.

Body Wt., lb.	Feed Consumption		Chlortetracycline		Laidlomycin	
	% Body Wt.	lb./head/day	Dose, mg/head/day	Feeding Rate, g/ton	Dose, mg/head/day	Feeding Rate g/ton
500	2.5	12.5	350	56	31.3	5
600	2	12	350	58.3	30	5
800	2.5	20	350	35	50	5
1000	3	30	350	23.3	75	5

Caution

Do not allow horses or other equines access to feeds containing laidlomycin propionate potassium. The safety of laidlomycin propionate potassium in unapproved species has not been established. Not for use in animals intended for breeding. If manufactured using liquid Type B laidlomycin propionate potassium feed, this dry Type C medicated feed expires 7 days after the date of manufacture.

Warning: No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

*Final printed label on formulated Type C Medicated feed must bear a single concentration for each drug.

^aAureomycin® is the proprietary name of chlortetracycline Type A medicated article (NADA 048-761).

^bCattlyst® is the proprietary name of laidlomycin propionate potassium Type A medicated article (NADA 141-025).

Zoetis Inc. [7 September 2016]