

**DCX + chlortetracycline-CMX Cattle Supplement Medicated  
Type C Medicated Feed**  
(decoquinatate and chlortetracycline Type C top-dress medicated feed)

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Calves, beef, and non-lactating dairy cattle. For the prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii*; for treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to chlortetracycline.

**ACTIVE DRUG INGREDIENTS\***

Decoquinatate <sup>a</sup> .....	90.9 to 535.7 g/ton*
Chlortetracycline <sup>b</sup> (as hydrochloride).....	4000 to 20,000 g/ton*

**GUARANTEED ANALYSIS**

Crude protein, not less than .....	_____ %
Crude fat, not less than .....	_____ %
Crude fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than .....	_____ %
Phosphorus, not less than.....	_____ %
Potassium, not less than.....	_____ %
Salt <sup>1</sup> , not less than.....	_____ %
Salt <sup>1</sup> , not more than .....	_____ %
Sodium <sup>2</sup> , not less than .....	_____ %
Sodium <sup>2</sup> , not more than .....	_____ %

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

**FEEDING DIRECTIONS**

Administer as a top dress supplement or mix into the daily ration to provide 22.7 mg decoquinatate per 100 lb of body weight per day and 1 g chlortetracycline per 100 lb body weight/day for not more than 5 days. \_\_\_\_\_\*\* pounds of this supplement will medicate 100 lbs of body weight. When it is fully consumed, resume feeding 22.7 mg decoquinatate per 100 lb of body weight/day for a total of 28 days to prevent coccidiosis.

**Caution:** Bentonite should not be used in decoquinatate feeds.

**Warning:** Withdraw 24 hours prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to animals producing milk for food.

Lot Number (if applicable): \_\_\_\_\_

**MANUFACTURED BY**  
Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

---

---

\*Final printed label on formulated Type C Medicated feed must bear a single concentration for each drug. The ratio between chlortetracycline and decoquinatone must be 44:1.

\*\*Actual concentration of chlortetracycline and decoquinatone will vary with body weight and feed consumption to provide 22.7 mg decoquinatone per 100 lb of body weight per day and 1 g chlortetracycline per 100 lb body weight/day for not more than 5 days. The following table provides examples of supplementation rates and amount of drug per ton of feed.

<b>Decoquinatone</b> Level in Supplement		<b>Chlortetracycline</b> Level in Supplement		Pounds of Supplement per 100 pounds of body weight
<b>mg/lb</b>	<b>grams/ton</b>	<b>mg/lb</b>	<b>grams/ton</b>	
45.45	90.9	2000	4000	0.5
56.75	113.5	2500	5000	0.4
113.5	227	5000	10,000	0.2
227	454	10,000	20,000	0.1

<sup>a</sup>Deccox® is the proprietary name of decoquinatone Type A medicated article (NADA 039-417).

<sup>b</sup>ChlorMax® is the proprietary name of chlortetracycline Type A medicated article (NADA 046-699).

Zoetis Inc. [15 September 2016]