

Veterinary Feed Directive for Use in Beef Cattle

Tylovet 100[®]

(tylosin phosphate)

Sequential VFD ID Number, if appropriate

Veterinarian: _____
 Address: _____
 Phone #: _____
 FAX or email: (optional) _____

Client: _____
 Business/Home Address: _____
 Phone #: _____
 FAX or email: (optional) _____

Indication, Drug Level in Medicated Feed, and Duration of Use: (specify additional required information)

For reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*.
Drug Level: _____ g/ton (8 to 10 g/ton)
Duration of Feeding: _____ days

USE OF FEED CONTAINING THIS VETERINARY FEED DIRECTIVE (VFD) DRUG IN A MANNER OTHER THAN AS DIRECTED ON THE LABELING (EXTRA-LABEL USE) IS NOT PERMITTED.

Approximate number of **Cattle** to be treated: _____
 Premise or Location of cattle: _____

Special Instructions and/or other animal identifications:

Affirmation of Intent (for combination VFD drugs): check the appropriate box:

- This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.
- This VFD authorizes the use of the VFD drug(s) cited in this order in the following FDA-approved, conditionally approved, or indexed combination(s) in medicated feed that contains the VFD drug(s) as a component.

	Drug(s) and Dose Range(s) [all listed doses are 90% DM basis]	Specifications*
<input type="checkbox"/>	Monensin at 10 to 40 g/ton to provide 0.14 to 0.42 mg/lb bodyweight/day Ractopamine at 9.8 to 24.6 g/ton to provide 90 to 430 mg/head/day (supplied by Rumensin [™] and Actogain [®] , ANADA 200-585)	For use in cattle fed in confinement for slaughter for the last 28 to 42 days on feed.
<input type="checkbox"/>	Monensin at 10 to 40 g/ton to provide 0.14 to 0.42 mg /lb bodyweight/day Ractopamine at 8.2 to 24.6 g/ton to provide 70 to 430 mg/head/day (supplied by Rumensin [™] and Actogain [®] , ANADA 200-585)	For use in cattle fed in confinement for slaughter for the last 28 to 42 days on feed.
<input type="checkbox"/>	Monensin at 10 to 40 g/ton to provide 0.14 to 0.42 mg/lb bodyweight/day Ractopamine up to 800 g/ton as a top dress to provide 70 to 400 mg/head/day (supplied by Rumensin [™] and Actogain [®] , ANADA 200-585)	For use in cattle fed in confinement for slaughter for the last 28 to 42 days on feed.
<input type="checkbox"/>	Monensin at 10 to 40 g/ton to provide 0.14 to 0.42 mg/lb bodyweight/day Ractopamine at 9.8 to 24.6 g/ton to provide 90 to 430 mg/head/day Melengestrol acetate, at the rate of 0.125 to 1.0 mg/lb (top dress or mixed at bunk with other Type C feed) to provide 0.25-0.5 mg/head/day. (supplied by Rumensin [™] and Actogain [®] and melengestrol acetate, MGA [®] , ANADA 200-583)	For use in cattle fed in confinement for slaughter for the last 28 to 42 days on feed.
<input type="checkbox"/>	Other FDA-approved, conditionally approved, or indexed combination:	

*for complete information see the approved Type C medicated feed label.

- This VFD authorizes the use of the VFD drug(s) cited in this order in any FDA-approved, conditionally approved, or indexed combination(s) in medicated feed that contains the VFD drug(s) as a component.

Warning: No withdrawal period is required.

Date of VFD Issuance: _____ (dd/mm/yyyy)

Date of VFD Expiration: _____ (dd/mm/yyyy)
(Cannot exceed 6 months after issuance)

Veterinarian's signature: _____

Color Z Original – Veterinarian
Color X Copy – Supplier
Color Y Copy – Client

All parties must retain a copy of this VFD for 2 years after issuance