

**DCX + Chlortetracycline-CMX Type B Cattle Feed Medicated
Type B Medicated Feed
(decoquinatate and chlortetracycline Type B medicated feed)**

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Calves, beef and non-lactating dairy cattle. For the prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii*; for treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENTS*

Decoquinatate^a 535.8 to 1818 g/ton
Chlortetracycline^b (as hydrochloride) 20,000 to 80,000 g/ton

GUARANTEED ANALYSIS

Crude protein, not less than %
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

MIXING DIRECTIONS

In order to treat cattle for bacterial enteritis and/or pneumonia while preventing coccidiosis, mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of complete cattle feed containing 12.9 to 90.8 grams decoquinatate and 500 to 4000 grams chlortetracycline, or a Type C medicated supplement containing 90.9 to 535.7 grams decoquinatate and 4000 to 20,000 grams chlortetracycline. The resulting Type C medicated feed should be fed at a rate to provide 1g chlortetracycline per 100 lb body weight/day and 22.7 mg decoquinatate per 100 lb of body weight/day daily for not more than 5 days. When this Type C medicated feed is fully consumed, resume feeding 22.7 mg decoquinatate per 100 lb of body weight/day for a total of 28 days to prevent coccidiosis. The following table provides examples of mixing rates.

Type B, grams/ton		Lbs Type B Per Ton of Feed	Lbs. Non-Medicated Feed Per Ton	Type C, grams/ton	
Decoquinatate	CTC			Decoquinatate	CTC
568	25,000	80	1920	22.7	1000
1136	50,000	40	1960	22.7	1000
1136	50,000	400	1600	227	10,000
1818	80,000	250	1750	227	10,000

Caution

Bentonite should not be used in decoquinatate feeds.

Warning: Withdraw 24 hours prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to animals producing milk for food.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

*Final printed label on formulated Type B Medicated feed must bear a single concentration for each drug. The ratio between chlortetracycline and decoquinatate must be 44:1.

^aDeccox® is the proprietary name of decoquinatate Type A medicated article (NADA 039-417).

^bChlorMax® is the proprietary name of chlortetracycline Type A medicated article (NADA 046-699).

Zoetis Inc. [15 September 2016]