

**Chlortetracycline-AUREO (350 mg/head) + Laidlomycin (5-10 g/ton)**  
**TYPE B Medicated Cattle Feed**  
(chlortetracycline and laidlomycin propionate potassium Type B medicated feed)

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline, and for improved feed efficiency in cattle fed in confinement for slaughter.

**ACTIVE DRUG INGREDIENTS\***

Chlortetracycline<sup>a</sup> ..... 127.3 to 4666.7 g/ton  
Laidlomycin propionate potassium<sup>b</sup> ..... 33.3 to 2000 g/ton

**GUARANTEED ANALYSIS**

Crude protein, not less than ..... %  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup> (not less than) ..... IU/lb.

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

**FEEDING DIRECTIONS**

In order to control bacterial pneumonia and improve feed efficiency, mix this Type B medicated feed with non-medicated feed ingredients to manufacture 1 ton of complete Type C medicated cattle feed containing 14.6 to 116.7 grams chlortetracycline and 5 to 10 grams laidlomycin propionate potassium. The following table provides examples of mixing rates.

Feed Consumption, lb./head/day	Type B Concentration, g/ton		Type B per ton Type C, lb.	Non-Medicated Feed per ton Type C, lb.	Type C Concentration, g/ton	
	Chlortetracyclin <sup>e</sup>	Laidlomycin			Chlortetracyclin <sup>e</sup>	Laidlomycin
12	388.9	33.3	300	1700	58.3	5
15	466.7	50.0	200	1800	46.7	5
20	700.0	150.0	100	1900	35.0	7.5
20	350.0	100.0	200	1800	35.0	10
30	155.6	50.0	300	1700	23.3	7.5
30	4666.7	2000.0	10	1990	23.3	10
36	3888.9	1500.0	10	1990	19.4	7.5

The resulting Type C medicated feed should be fed continuously at a rate to provide 350 mg chlortetracycline per head per day and 30 to 150 mg laidlomycin propionate potassium per head per day.

**Caution**

Do not allow horses or other equines access to feeds containing laidlomycin propionate potassium. The safety of laidlomycin propionate potassium in unapproved species has not been established. Not for use in animals intended for breeding.

**Warning:** No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Lot Number (if applicable): \_\_\_\_\_

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

\*Final printed label on formulated Type C Medicated feed must bear a single concentration for each drug.

<sup>a</sup>Aureomycin® is the proprietary name of chlortetracycline Type A medicated article (NADA 048-761).

<sup>b</sup>Cattlyst® is the proprietary name of laidlomycin propionate potassium Type A medicated article (NADA 141-025).

Zoetis Inc. [13 June 2016]