Veterinary Feed Directive for Turkeys Aureomycin[®] (chlortetracycline)

Veterinariar	n:		
Address:		Business or Home Address:	
	ail: (optional)	Phone #:	
	, Drug Level in Medicated Fee	ed, and Duration of Use: (select one and spec	
	Turkeys: Control of compli (transmissible enteritis, co	icating bacterial organisms associated w ronaviral enteritis) susceptible to chlorte to provide 25 mg/lb body weight/day) lays (7 to 14 days)	
2)	Turkeys: Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. Drug level: 200 g/ton Duration of use: days (7 to 14 days)		
3)	Turkeys: Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to chlortetracycline. Drug level: 400 g/ton Duration of use: days (7 to 14 days)		
4)		veeks of age: Reduction of mortality due n susceptible to chlortetracycline. days	to paratyphoid caused
	FEED CONTAINING THIS VET	ERINARY FEED DIRECTIVE (VFD) DRUG IN LABELING (EXTRA-LABEL USE) IS NOT PI	A MANNER OTHER THAN AS
••	-	ated:	
Premise o	r Location of animals:		
Special Ins	structions and/or other animal id	entifications:]
Affirmatio	on of Intent (for combination VF	D drugs):	
	only authorizes the use of the VF combination with any other anim	D drug(s) cited in this order and is not intended nal drugs.	to authorize the use of such
	Warning: No withdrawa	al period is required. Do not feed to turkeys pro for human consumption.	ducing eggs
Date of VF	D Issuance:(dd/mm/yyyy) Date of VFD Expiration:_ (Cannot exceed 6 months a	
Veterinarian	s signature: Color Z Original – Veterinarian	Color X Copy – Supplier	Color Y Copy – Client
		s must retain a copy of this VFD for 2 years after issuance	oolor i oopy – onent