## (sulfadimethoxine/ormethoprim)

Veterinary Feed Directive for Ducks			
RofenAid <sup>®</sup>			

Veterinarian:		Business or
Address:		Home Address:
	ail: (optional)	Phone #: FAX or email: (optional)
	s, Drug Level, and Duration of Use: (specify	
	<b>Drug level:</b> 227 g sulfadimethoxine and 13 <b>Duration of use:</b> 7 days	6.2 g ormethoprim per ton
2)	<b>Ducks:</b> As an aid in the control of <i>Escherica</i> severe challenge of <i>P. multocida</i> (fowl choles)	hia coli, Riemerella anatipestifer (strains 2, 3 & 4) and era).
	<b>Drug level:</b> 454 g sulfadimethoxine and 27 <b>Duration of use:</b> 7 days	2.4 g ormethoprim per ton
USE OF	FEED CONTAINING THIS VETERINARY F	EED DIRECTIVE (VFD) DRUG IN A MANNER OTHER THAN AS
	DIRECTED ON THE LABELING	6 (EXTRA-LABEL USE) IS NOT PERMITTED.
Premise c	or Location of animals:	
Special In	structions and/or other animal identifications:	
Affirmation	on of Intent (for combination VFD drugs): ch	eck the appropriate box:
	only authorizes the use of the VFD drug(s) cin combination with any other animal drugs.	ited in this order and is not intended to authorize the use of such
Fo	Medication should be started at the first sign r indication 2 only: Not for breeding ducks. her than 0.025% and 0.015%, respectively, i	The safety of feeding sulfadimethoxine and ormetoprim at levels
		raw 5 days before slaughter. cks producing eggs for food.
Date of VF	D Issuance:(dd/mm/yyyy)	Date of VFD Expiration: (dd/mm/yyyy) (Cannot exceed 6 months after issuance)
√eterinaria	n's signature:	