Veterinary Feed Directive for Turkeys RofenAid®

(Sunadimethoxine/ormethoprim)	
√eterinarian:	Client:
	Business or
Address:	
Phone #: FAX or email: (optional)	FAX or email: (optional)
ndications, Drug Level, and Duration of Use	
-	
Furkeys: As an aid in the prevention of coccid for the prevention of bacterial infections cause	iosis by Eimeria adenoeides, E. gallopavonis, E. meleagrimitis;
or the prevention of bacterial infections caused	a by Pasteurella Multocida (IOWI Cholera).
Orug level: 56.75 g sulfadimethoxine and 34.0	95 g ormethoprim per ton
Donathan afaire	
Duration of use:days	***************************************
	INARY FEED DIRECTIVE (VFD) DRUG IN A MANNER OTHER THAN AS
	ABELING (EXTRA-LABEL USE) IS NOT PERMITTED.
	d:
Premise or Location of animals:	
Special Instructions and/or other animal identi	ifications:
opecial instructions and/or other animal identity	incations.
Affirmation of Intent (for combination VFD d	ruas):
•	drug(s) cited in this order and is not intended to authorize the use of such
drug(s) in combination with any other animal	
	eed to turkeys producing eggs for food.
Do not le	to tarkeys producing eggs for food.
Date of VFD Issuance:(dd/mm/yyyy)	Date of VFD Expiration: (dd/mm/yyyy)
	(Cannot exceed 6 months after issuance)

Color Z Original – Veterinarian

Veterinarian's signature:

Color Y Copy - Client

harian Color X Copy – Supplier
All parties must retain a copy of this VFD for 2 years after issuance