

Lincomycin SWINE RATION L100
Type C Medicated Complete Feed
(lincomycin Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For treatment of swine dysentery and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*.

ACTIVE DRUG INGREDIENT

Lincomycin^a (as lincomycin hydrochloride agricultural grade).....100 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Lysine, not less than.....	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%
Selenium, not less than	_____	ppm
Zinc, not less than	_____	ppm

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be specifically named in accordance with the names and definitions adopted by the AAFCO.

DIRECTIONS FOR USE

Feed 100 grams of lincomycin per ton of complete feed as the sole ration for three weeks or until clinical signs of the disease (watery, mucoid, or bloody stools) disappear.

CAUTION

Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment. The effects of lincomycin on swine reproductive performance, pregnancy, and lactation have not been determined.

Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.



WARNING: No drug withdrawal period is required before slaughter of swine fed lincomycin at 100 grams lincomycin per ton of feed.



To report adverse effects, access medical information, or obtain additional product information, call 1-888-963-8471.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

^aLincomix® is the proprietary name of lincomycin formulation Type A medicated article (NADA 097-505).

Zoetis Inc. [14 March 2016]