

Lincomycin BROILER Feed
Lincomycin Type C Medicated Complete Feed
(lincomycin Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the control of necrotic enteritis caused or complicated by *Clostridium spp.* or other organisms susceptible to lincomycin in broilers.

ACTIVE DRUG INGREDIENT

Lincomycin^a (as lincomycin hydrochloride agricultural grade).....2 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Lysine, not less than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than	_____	%
Salt ¹ , not less than	_____	%
Salt ¹ , not more than	_____	%
Sodium ² , not less than	_____	%
Sodium ² , not more than	_____	%
Selenium, not less than.....	_____	ppm
Zinc, not less than.....	_____	ppm

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be specifically named in accordance with the names and definitions adopted by the AAFCO.

DIRECTIONS FOR USE

Feed as the sole ration to broilers.

CAUTION

Not for use in layers, breeders, or turkeys.

Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.



WARNING: No drug withdrawal period is required before slaughter of birds fed lincomycin at 2 grams lincomycin per ton of feed.



To report adverse effects, access medical information, or obtain additional product information, call 1-888-963-8471.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

^aLincomix® is the proprietary name of lincomycin formulation Type A medicated article (NADA 097-505).

Zoetis Inc. [14 March 2016]