

# Veterinary Feed Directive for Swine Aureomycin® (chlortetracycline)

Veterinarian: \_\_\_\_\_  
Address: \_\_\_\_\_  
Phone #: \_\_\_\_\_  
FAX or email: (optional) \_\_\_\_\_

Client: \_\_\_\_\_  
Business or Home Address: \_\_\_\_\_  
Phone #: \_\_\_\_\_  
FAX or email: (optional) \_\_\_\_\_

\*\*\*\*\*  
**Indications, Drug Level in Medicated Feed, and Duration of Use: (select one and specify additional required information)**

- 1) Swine: Control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis* susceptible to chlortetracycline.  
**Drug Concentration:** \_\_\_\_\_g/ton (to provide 10 mg/lb body weight/day, which is equivalent to approximately 400 g/ton)  
**Duration of use:** \_\_\_\_\_days (Feed for not more than 14 days)
- 2) Swine: Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to chlortetracycline.  
**Drug Concentration:** \_\_\_\_\_g/ton (to provide 10 mg/lb body weight/day, which is equivalent to approximately 400 g/ton)  
**Duration of use:** \_\_\_\_\_days (feed for not more than 14 days)
- 3) Swine: Reduction in the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E *Streptococci* susceptible to chlortetracycline.  
**Drug Concentration:** \_\_\_\_\_g/ton (50 to 100 g/ton)  
**Duration of use:** \_\_\_\_\_days
- 4) Breeding Swine: Control of leptospirosis (reducing the incidence of abortion and shedding of *leptospirea*) caused by *Leptospira pomona* susceptible to chlortetracycline.  
**Drug Concentration:** 400 g/ton  
**Duration of use:** \_\_\_\_\_days (feed continuously for not more than 14 days)

\*\*\*\*\*  
**USE OF FEED CONTAINING THIS VETERINARY FEED DIRECTIVE (VFD) DRUG IN A MANNER OTHER THAN AS DIRECTED ON THE LABELING (EXTRA-LABEL USE) IS NOT PERMITTED.**

Approximate number of **Swine** to be treated: \_\_\_\_\_  
Premises or Location of animals: \_\_\_\_\_

Special Instructions and/or other animal identifications:

**Affirmation of Intent (for combination VFD drugs): check the appropriate box:**

- This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.
- This VFD authorizes the use of the VFD drug(s) cited in this order in the following FDA-approved, conditionally approved, or indexed combination(s) in medicated feed that contains the VFD drug(s) as a component.

	Drug(s) and Dose Range(s)	Specifications*
<input type="checkbox"/>	10 to 30 g/ton bacitracin methylene disalicylate; (BMD®) [NADA 200-242]	Growing/finishing swine, pregnant sows This combination has a zero day withdrawal.
<input type="checkbox"/>	Other FDA-approved, conditionally approved, or indexed combination:	

\*for complete information see the approved Type C medicated feed label

- This VFD authorizes the use of the VFD drug(s) cited in this order in any FDA-approved, conditionally approved, or indexed combination(s) in medicated feed that contains the VFD drug(s) as a component.

**Withdrawal Period:** No withdrawal period is required.

Date of VFD Issuance: \_\_\_\_\_ (dd/mm/yyyy)

Date of VFD Expiration: \_\_\_\_\_ (dd/mm/yyyy)  
(Cannot exceed 6 months after issuance)

Veterinarian's signature: \_\_\_\_\_

**Color Z Original – Veterinarian**

**Color X Copy – Supplier**

**Color Y Copy – Client**

All parties must retain a copy of this VFD for 2 years after issuance