Veterinary Feed Directive for Use in Swine Tylovet 100® (tylosin phosphate)

Sequential VFD ID Number, if appropriate

/otorinarian	Client:			
AX or email:	(optional)	Phone #: FAX or email: (optional)		
	Drug Level in Medicated Feed	, and Duration of Use: (select one and specify additional required		
ш	For reduction in severity of effe Drug Level: 100 grams per tor Duration of Feeding:			
	For control of swine dysentery a Drug Level: 100 grams per tor Duration of Feeding:			
	Subsequent Drug Level: 40 g Duration of Feeding:	rams per ton _weeks (until pigs reach market weight)		
	For the treatment and control of swine dysentery associated with <i>Brachyspira hyodysenteria</i> immediately after medicating with tylosin tartrate in drinking water. Drug Level:grams per ton (40 to 100 g/ton) Duration of Feeding:weeks (2 to 6 weeks)			
	For control of porcine proliferative enteropathies (PPE, ileitis) associated with <i>Lawsonia intracellularis</i> . Drug Level: 100 grams per ton Duration of Feeding: 21 days			
	For control of porcine proliferative enteropathies (PPE, ileitis) associated with <i>Lawsonia intracellularis</i> . Drug Level: 100 grams per ton Duration of Feeding: weeks (minimum of 3 weeks)			
	Subsequent Drug Level: 40 g Duration of Feeding:	rams per ton _weeks (until pigs reach market weight)		
	For control of porcine proliferative enteropathies (PPE, ileitis) associated with <i>Lawsonia intracellularis</i> immediately after medicating with tylosin tartrate in drinking water. Drug Level: grams per ton (40 to 100 g/ton) Duration of Feeding: weeks (2 to 6 weeks)			

Page 1 of 2

*****	*****	********************	*************		
		USE OF FEED CONTAINING THIS VETERINARY FEED DIRECTIVE (VF DIRECTED ON THE LABELING (EXTRA-LABEL US			
Approxi	imate n	number of Swine to be treated:			
		cation of swine:			
		ctions and/or other animal identifications:			
		f Intent (for combination VFD drugs): check the appropriate box:			
	This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.				
	This VFD authorizes the use of the VFD drug(s) cited in this order in the following FDA-approved, conditionally approved, or				
_		ked combination(s) in medicated feed that contains the VFD drug(s)	as a component.		
		Drug(s) and Dose Range(s)	Specifications*		
		ractopamine hydrochloride at 4.5 to 9.0 g/ton** (supplied by	For use in finishing swine weighing not less		
		Engain®, ANADA 200-584)	than 150 lbs for the last 45 to 90 lbs (group		
		Other FDA-approved, conditionally approved, or indexed	average) of weight gain prior to slaughter.		
		combination:			
		*for complete information see the approved Type C medicated fe	ed label		
		**combination not available with atrophic rhinitis indication			
		VFD authorizes the use of the VFD drug(s) cited in this order in any pination(s) in medicated feed that contains the VFD drug(s) as a con			
	COITIL	ornation(s) in medicated feed that contains the VFD drug(s) as a con-	iiponent.		
		Warning: No withdrawal period is	required.		
Data of \	/ED les	suance: (dd/mm/yyyy) Date of V	FD Expiration: (dd/mm/yyyy)		
Date of v	/1 10 133		ceed 6 months after issuance)		
Veterina	arian's	signature:	,		
	C	Color Z Original – Veterinarian Color X Copy – Supplie			
		All parties must retain a copy of this VFD for 2 year	rs after issuance		

Page 2 of 2