

Sulfadimethoxine and Ormetoprim Chicken Feed^a
Type C Medicated Complete Feed
(sulfadimethoxine and ormetoprim Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Broiler and Replacement Chickens (Breeder and Layer): As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*; as an aid in the prevention of bacterial infections caused by *Haemophilus gallinarum* (infectious coryza), *Escherichia coli* (colibacillosis) and *Pasteurella multocida* (fowl cholera).

Active Drug Ingredients^a

Sulfadimethoxine.....	113.5 g/ton (0.0125%)
Ormetoprim.....	68.1 g/ton (0.0075%)

GUARANTEED ANALYSIS

Crude Protein, Min	%
Lysine, Min.....	%
Methionine, Min	%
Crude Fat, Min.....	%
Crude Fiber, Max.....	%
Calcium, Min	%
Calcium ¹ Max	%
Phosphorus, Min.....	%
Salt ¹ , Min.....	%
Salt ¹ , Max	%
Sodium ² , Min	%
Sodium ² , Max.....	%

¹ Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

² Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Ingredients as defined by AAFCO

Directions for Use

Feed continuously as the sole ration.

Warning

Withdraw 5 days before slaughter. Do not feed to chickens over 16 weeks (112 days) of age.

Lot Number (if applicable):_____

MANUFACTURED BY

Blue Bird Feed Company

City, State, Zip

NET WEIGHT ON BAG OR BULK

^aRofenAid® 40 is the proprietary name of sulfadimethoxine and ormetoprim formulation Type A medicated article (NADA 040-209).