

Chlortetracycline Turkey Feed – BLUECOMB - CMX^a
Type C Medicated Complete Feed
(chlortetracycline Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline.

Active Drug Ingredient

Chlortetracycline^a (____g/ton)**

** The final printed feed label must show actual concentration of chlortetracycline, which will vary with body weight and feed consumption to provide 25 mg/lb body weight per day.

Guaranteed Analysis

Crude Protein (Min).....	_____	%
Lysine (Min).....	_____	%
Methionine (Min).....	_____	%
Crude Fat (Min)	_____	%
Crude Fiber (Max)	_____	%
Calcium (Min).....	_____	%
Calcium (Max).....	_____	%
Phosphorus (Min)	_____	%
Salt ¹ (Min)	_____	%
Salt ¹ (Max).....	_____	%
Sodium ² (Min).....	_____	%
Sodium ² (Max)	_____	%

¹ Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

² Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Ingredients as defined by AAFCO.

Feeding Directions

Feed continuously for 7 to 14 days, providing chlortetracycline at the rate of 25 mg per pound of body weight per day. The following table provides examples of feeding rates and amount of drug in feed.

Drug Level in Feed		Lbs of Turkey Feed Per 100 lbs Body Weight
mg/lb	grams/ton	
500	1000	5.0
625	1250	4.0
833	1667	3.0
1250	2500	2.0

Warning

Do not feed to turkeys producing eggs for human consumption.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

^aChlorMax® 50 is the proprietary name of chlortetracycline Type A medicated article.

Zoetis Inc. [20 January 2016]