

**Chlortetracycline Turkey Feed – AUREO<sup>a</sup> - SYN  
Type B Medicated Feed**  
(chlortetracycline Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline in turkeys.

**Active Drug Ingredient**

Chlortetracycline<sup>a</sup> ..... 201 – 80,000 g/ton\*

**Guaranteed Analysis**

|                                 |   |
|---------------------------------|---|
| Crude Protein (Min) .....       | % |
| Lysine (Min).....               | % |
| Methionine (Min) .....          | % |
| Crude Fat (Min).....            | % |
| Crude Fiber (Max).....          | % |
| Calcium (Min) .....             | % |
| Calcium (Max).....              | % |
| Phosphorus (Min) .....          | % |
| Salt <sup>1</sup> (Min).....    | % |
| Salt <sup>1</sup> (Max).....    | % |
| Sodium <sup>2</sup> (Min) ..... | % |
| Sodium <sup>2</sup> (Max).....  | % |

<sup>1</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>2</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Ingredients as defined by AAFCO.

**Mixing Directions**

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for turkeys containing 200 grams chlortetracycline per ton.

*The following table provides examples of mixing rates:*

| Type B CTC concentration (g/ton) | Type B per ton of Type C (lb) | Non-medicated feed per ton of Type C (lb) | Type C CTC concentration (g/ton) |
|----------------------------------|-------------------------------|---|----------------------------------|
| 10,000 (5 g/lb)                  | 40                            | 1960                                      | 200                              |
| 20,000 (10 g/lb)                 | 20                            | 1980                                      | 200                              |
| 40,000 (20 g/lb)                 | 10                            | 1990                                      | 200                              |
| 80,000 (40 g/lb)                 | 5                             | 1995                                      | 200                              |

The resulting Type C medicated feed should be fed continuously as the sole ration for 7 to 14 days.

**Warning**

Do not feed to turkeys producing eggs for human consumption.

Lot Number (if applicable):\_\_\_\_\_

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

\*The final printed medicated feed label must state a single drug concentration.

<sup>a</sup>Aureomycin® is the proprietary name of chlortetracycline Type A medicated article (NADA 048-761).

Zoetis Inc. [15 March 2016]