

**Lincomycin/Pyrantel SWINE RATION LB-MP**  
**Type C Medicated Complete Feed**  
(lincomycin/pyrantel tartrate Type C medicated feed)

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For reduction in the severity of mycoplasmal pneumonia caused by Mycoplasma hyopneumoniae. Aid in the prevention of migration and establishment of large roundworm (Ascaris suum) infections; aid in the prevention of establishment of nodular worm (Oesophagostomum spp) infections of swine when fed in accordance with directions for use.

**ACTIVE DRUG INGREDIENTS**

Lincomycin (as lincomycin hydrochloride agricultural grade)<sup>a</sup> .....200 g/ton  
(Each pound contains 0.1 gram lincomycin)  
Pyrantel tartrate<sup>b</sup>.....0.0106% (96 g/ton)  
(Each pound contains 0.048 grams of pyrantel tartrate)

**GUARANTEED ANALYSIS**

Crude Protein, not less than..... %  
Crude Fat, not less than..... %  
Crude Fiber, not more than..... %

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. Include also ingredients of the trace mineral and vitamin premixes.

**CAUTION**

Occasionally, swine fed lincomycin may within the first two days after the onset of treatment development diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self correcting within five to eight days without discontinuing the lincomycin treatment.

The effects of lincomycin on swine reproductive performance, pregnancy, and lactation have not been determined.

Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

Do not mix in feeds containing bentonite.

Not for use in swine that weigh more than 250 pounds.



**WARNING:** Do not slaughter swine for human consumption for 6 days following last treatment.



**DIRECTIONS FOR USE**

Feed 200 grams of lincomycin and 96 grams of pyrantel tartrate per ton of complete feed as the sole ration for 21 days.

Lot Number (if applicable): \_\_\_\_\_

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

<sup>a</sup>Lincomix® is the proprietary name of lincomycin formulation Type A medicated article (NADA 097-505).

<sup>b</sup>Banminth® is the proprietary name of pyrantel tartrate formulation Type A medicated article (NADA 043-290).