

**Chlortetracycline - CMX/Monensin Chicken Feed  
Type C Medicated Complete Feed  
(chlortetracycline/monensin Type C medicated feed)**

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

As an aid in the reduction of mortality due to *E. coli* infections susceptible to chlortetracycline in chickens.  
As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

**ACTIVE DRUG INGREDIENTS**

Chlortetracycline<sup>a</sup> (as HCl).....500 g/ton  
Monensin<sup>b</sup>.....90 to 110 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein (Min).....	_____	%
Lysine (Min).....	_____	%
Methionine (Min).....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium (Min).....	_____	%
Calcium (Max).....	_____	%
Phosphorus (Min).....	_____	%
Salt <sup>1</sup> (Min).....	_____	%
Salt <sup>1</sup> (Max).....	_____	%
Sodium <sup>2</sup> (Min).....	_____	%
Sodium <sup>2</sup> (Max).....	_____	%

<sup>1</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>2</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**DIRECTIONS FOR USE**

Feed for five days as the sole ration.

**CAUTION:** Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. In the absence of coccidiosis in broiler chickens, use of monensin with no withdrawal period may limit feed intake, resulting in reduced gains. Monensin is approved for use in replacement chickens for use as cage layers only.



**WARNING:** Withdraw 24 hours before slaughter. Do not feed to laying chickens.

Not to be fed continuously for more than 5 days.

Do not feed to chickens over 16 weeks of age.



Lot Number (if applicable): \_\_\_\_\_

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

\*Final Type C medicated feed must list a single level of monensin.

<sup>a</sup>ChlorMax® is the proprietary name of chlortetracycline Type A medicated article (NADA 046-699).

<sup>b</sup>Coban® is the proprietary name of monensin Type A medicated article (NADA 038-878).

Zoetis Inc. [14 November 2016]