

Veterinary Feed Directive for Chickens ChlorMax® (chlortetracycline)

Veterinarian: _____

Client: _____

Address: _____

Business or Home Address: _____

Phone #: _____

Phone #: _____

FAX or email: (optional) _____

FAX or email: (optional) _____

Indications, Drug Level, and Duration of Use: (select one and specify additional required information)

1) Chickens: Control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline.

Drug level: _____ g/ton (100 to 200 g/ton)

Duration of use: _____ days (7 to 14 days)

2) Chickens: Control of chronic respiratory disease (CRD) and air sac infection caused by *Mycoplasma gallisepticum* and *Escherichia coli* susceptible to chlortetracycline.

Drug level: _____ g/ton (200 to 400 g/ton)

Duration of use: _____ days (7 to 14 days)

3) Chickens: Reduction of mortality due to *Escherichia coli* infections susceptible to chlortetracycline.

Drug level: 500 g/ton

Duration of use: 5 days

USE OF FEED CONTAINING THIS VETERINARY FEED DIRECTIVE (VFD) DRUG IN A MANNER OTHER THAN AS DIRECTED ON THE LABELING (EXTRA-LABEL USE) IS NOT PERMITTED.

Approximate number of **Chickens** to be treated: _____

Premises or Location of animals: _____

Special Instructions and/or other animal identifications:

Affirmation of Intent (for combination VFD drugs): check the appropriate box:

This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.

This VFD authorizes the use of the VFD drug(s) cited in this order in the following FDA-approved, conditionally approved, or indexed combination(s) in medicated feed that contains the VFD drug(s) as a component.

	Drug(s) and Dose Range(s)	Specifications*
<input type="checkbox"/>	90 to 110 g/ton monensin (COBAN®) [ANADA 200-263]	For use in broiler chickens. Not to be fed continuously for more than 5 days. Withdraw 24 hours before slaughter.
<input type="checkbox"/>	40 to 60 g/ton salinomycin (BIOCOX®) [ANADA 200-261]	For use in broiler chickens. Not to be fed for more than 5 days. Withdraw 24 hours before slaughter.
<input type="checkbox"/>	40 to 60 g/ton salinomycin (SACOX®) [ANADA 200-262]	For use in broiler chickens. Not to be fed for more than 5 days. Withdraw 24 hours before slaughter.
<input type="checkbox"/>	Other FDA-approved, conditionally approved, or indexed combination:	

*for complete information see the approved Type C medicated feed label

This VFD authorizes the use of the VFD drug(s) cited in this order in any FDA-approved, conditionally approved, or indexed combination(s) in medicated feed that contains the VFD drug(s) as a component.

Warning: Do not feed to chickens producing eggs for human consumption.

Indication 1, 2: No withdrawal period required.

Indication 3: Withdraw 24 hours prior to slaughter.



Date of VFD Issuance: _____(dd/mm/yyyy)

Date of VFD Expiration: _____ (dd/mm/yyyy)
(Cannot exceed 6 months after issuance)

Veterinarian's signature: _____

Color Z Original – Veterinarian

Color X Copy – Supplier

Color Y Copy – Client

All parties must retain a copy of this VFD for 2 years after issuance