

Veterinary Feed Directive for Swine Lincomix® (lincomycin)

Veterinarian: _____
Address: _____
Phone #: _____
FAX or email: (optional) _____

Client: _____
Business or Home Address: _____
Phone #: _____
FAX or email: (optional) _____

Indications, Drug Level, and Duration of Use: (select one and specify additional required information)

- 1) For reduction in the severity of swine mycoplasmal pneumonia caused by *Mycoplasma hyopneumoniae*.
Drug level: 200 g/ton
Duration of use: 21 days

USE OF FEED CONTAINING THIS VETERINARY FEED DIRECTIVE (VFD) DRUG IN A MANNER OTHER THAN AS DIRECTED ON THE LABELING (EXTRA-LABEL USE) IS NOT PERMITTED.

Approximate number of **Swine** to be treated: _____
Premises or Location of animals: _____
Special Instructions and/or other animal identifications:

Affirmation of Intent (for combination VFD drugs): **check the appropriate box:**

- This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.
- This VFD authorizes the use of the VFD drug(s) cited in this order in the following FDA-approved, conditionally approved, or indexed combination(s) in medicated feed that contains the VFD drug(s) as a component.

Drug(s) and Dose Range(s)	Specifications*
<input type="checkbox"/> Pyrantel tartrate – 96 g/ton in Type C medicated feed – (BANMINTH®) [NADA 138-941]	Withdraw 6 days before slaughter Not to be fed to swine weighing more than 250 lb.
<input type="checkbox"/> Other FDA-approved, conditionally approved, or indexed combination:	

*for complete information see the approved Type C medicated feed label.

- This VFD authorizes the use of the VFD drug(s) cited in this order in any FDA-approved, conditionally approved, or indexed combination(s) in medicated feed that contains the VFD drug(s) as a component.

▶ **WARNING:** When using LINCOMIX 20 or LINCOMIX 50 in approved combinations with other drugs, follow the required withdrawal times for those drugs. No drug withdrawal period is required before slaughter of swine fed LINCOMIX 20 or LINCOMIX 50 at approved concentrations (40, 100 or 200 grams lincomycin per ton of feed). ◀

CAUTION: Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions are self-correcting within five to eight days without discontinuing the lincomycin treatment.

The effects of lincomycin on swine reproductive performance, pregnancy, and lactation have not been determined.

Do not allow rabbits, guinea pigs, horses or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

Date of VFD Issuance: _____ (dd/mm/yyyy) Date of VFD Expiration: _____ (dd/mm/yyyy)
(Cannot exceed 6 months after issuance)

Veterinarian's signature: _____